

### Decamp Study 1: Diagnosis and Surveillance of Indeterminate Pulmonary Nodules

#### Inclusion Criteria

- Patients aged 45 years or older
- Diagnosed with an indeterminate pulmonary nodule (0.7 - 3.0 cm)
- Nodules must be of appropriate size at enrollment, but nodule(s) may have been first identified within 12 months prior to enrollment
- Current or former smoker with  $\geq 20$  pack years (pack years = number of packs smoked per day x number of years smoked)
- Willing to undergo fiberoptic bronchoscopy
- Able to tolerate biospecimen collections
- Able to comply with standard of care (initial and follow up) visits including clinical exams, diagnostic work-ups, and imaging for two years

#### Exclusion Criteria

- History or previous diagnosis of lung cancer
- History of diagnoses of pure ground glass opacities w/ no indeterminate pulmonary nodules
- Contraindications to nasal brushing or fiberoptic bronchoscopy procedure
- Allergies to any local anesthetic that may be used to obtain biosamples in the study

If you have a patient who may qualify for this study, please contact:

Version: 08-06-2014

### Decamp Study 1: Diagnosis and Surveillance of Indeterminate Pulmonary Nodules

#### Inclusion Criteria

- Patients aged 45 years or older
- Diagnosed with an indeterminate pulmonary nodule (0.7 - 3.0 cm)
- Nodules must be of appropriate size at enrollment, but nodule(s) may have been first identified within 12 months prior to enrollment
- Current or former smoker with  $\geq 20$  pack years (pack years = number of packs smoked per day x number of years smoked)
- Willing to undergo fiberoptic bronchoscopy
- Able to tolerate biospecimen collections
- Able to comply with standard of care (initial and follow up) visits including clinical exams, diagnostic work-ups, and imaging for two years

#### Exclusion Criteria

- History or previous diagnosis of lung cancer
- History of diagnoses of pure ground glass opacities w/ no indeterminate pulmonary nodules
- Contraindications to nasal brushing or fiberoptic bronchoscopy procedure
- Allergies to any local anesthetic that may be used to obtain biosamples in the study

If you have a patient who may qualify for this study, please contact:

Version: 08-06-2014

### Decamp Study 1: Diagnosis and Surveillance of Indeterminate Pulmonary Nodules

#### Inclusion Criteria

- Patients aged 45 years or older
- Diagnosed with an indeterminate pulmonary nodule (0.7 - 3.0 cm)
- Nodules must be of appropriate size at enrollment, but nodule(s) may have been first identified within 12 months prior to enrollment
- Current or former smoker with  $\geq 20$  pack years (pack years = number of packs smoked per day x number of years smoked)
- Willing to undergo fiberoptic bronchoscopy
- Able to tolerate biospecimen collections
- Able to comply with standard of care (initial and follow up) visits including clinical exams, diagnostic work-ups, and imaging for two years

#### Exclusion Criteria

- History or previous diagnosis of lung cancer
- History of diagnoses of pure ground glass opacities w/ no indeterminate pulmonary nodules
- Contraindications to nasal brushing or fiberoptic bronchoscopy procedure
- Allergies to any local anesthetic that may be used to obtain biosamples in the study

If you have a patient who may qualify for this study, please contact:

Version: 08-06-2014

### Decamp Study 1: Diagnosis and Surveillance of Indeterminate Pulmonary Nodules

#### Inclusion Criteria

- Patients aged 45 years or older
- Diagnosed with an indeterminate pulmonary nodule (0.7 - 3.0 cm)
- Nodules must be of appropriate size at enrollment, but nodule(s) may have been first identified within 12 months prior to enrollment
- Current or former smoker with  $\geq 20$  pack years (pack years = number of packs smoked per day x number of years smoked)
- Willing to undergo fiberoptic bronchoscopy
- Able to tolerate biospecimen collections
- Able to comply with standard of care (initial and follow up) visits including clinical exams, diagnostic work-ups, and imaging for two years

#### Exclusion Criteria

- History or previous diagnosis of lung cancer
- History of diagnoses of pure ground glass opacities w/ no indeterminate pulmonary nodules
- Contraindications to nasal brushing or fiberoptic bronchoscopy procedure
- Allergies to any local anesthetic that may be used to obtain biosamples in the study

If you have a patient who may qualify for this study, please contact:

Version: 08-06-2014