










# Radiation Dose to Adults From Common Imaging Examinations

Procedure		Approximate effective radiation dose (mSv)	Approximate comparable time of natural background radiation exposure
 <b>ABDOMINAL REGION</b>	Computed Tomography (CT) — Abdomen and Pelvis	7.7 mSv	2.6 years
	Computed Tomography (CT) — Abdomen and Pelvis, repeated with and without contrast material	15.4 mSv	5.1 years
	Computed Tomography (CT) — Colonography	6 mSv	2 years
	Intravenous Urogram (IVU))	3 mSv	1 year
	Barium Enema (Lower GI X-ray)	6 mSv	2 years
	Upper GI Study With Barium	6 mSv	2 years
 <b>BONE</b>	Lumbar Spine	1.4 mSv	6 months
	Extremity (hand, foot, etc.) X-ray	< 0.001 mSv	< 3 hours
 <b>CENTRAL NERVOUS SYSTEM</b>	Computed Tomography (CT) — Brain	1.6 mSv	7 months
	Computed Tomography (CT) — Brain, repeated with and without contrast material	3.2 mSv	13 months
	Computed Tomography (CT) — Head and Neck	1.2 mSv	5 months
	Computed Tomography (CT) — Spine	8.8 mSv	3 years
 <b>CHEST</b>	Computed Tomography (CT) — Chest	6.1 mSv	2 years
	Computed Tomography (CT) — Lung Cancer Screening	1.5 mSv	6 months
	Chest X-ray	0.1 mSv	10 days
 <b>DENTAL</b>	Dental X-ray	0.005 mSv	1 day
	Panoramic X-Ray	0.025 mSv	3 days
	Cone Beam CT	0.18 mSv	22 days
 <b>HEART</b>	Coronary Computed Tomography Angiography (CTA)	8.7 mSv	3 years
	Cardiac CT for Calcium Scoring	1.7 mSv	6 months
	Non-Cardiac Computed Tomography Angiography (CTA)	5.1 mSv	< 2 years
 <b>MEN'S IMAGING</b>	Bone Densitometry (DEXA)	0.001 mSv	3 hours
 <b>NUCLEAR MEDICINE</b>	Positron Emission Tomography — Computed Tomography (PET/CT) Whole body protocol	22.7 mSv	7.6 years
 <b>WOMEN'S IMAGING</b>	Bone Densitometry (DEXA)	0.001 mSv	3 hours
	Screening Digital Mammography	0.21 mSv	26 days
	Screening Digital Breast Tomosynthesis (3D Mammogram)	0.27 mSv	33 days

**Note:** This chart simplifies a highly complex topic for patients' informational use. The effective doses are typical values for an average-sized adult. The actual dose can vary substantially, depending on a person's size as well as on differences in imaging practices. It is also important to note that doses given to pediatric patients will vary significantly from those given to adults, since children vary in size. Patients with radiation dose questions should consult with their medical physicists and/or radiologists as part of a larger discussion on the benefits and risks of radiologic care.